

SUR L'ESTIMATION DES ERREURS EN
CONVERGENCE NUMÉRIQUE DE CERTAINES
MÉTHODES D'ITÉRATION

(English translation)

ON THE ESTIMATION OF ERRORS IN NUMERICAL
CONVERGENCE OF CERTAIN ITERATION METHODS

ION PĂVĂLOIU

Let E be a Banach space and

$$(1) \quad x = \lambda D(x) + y, \quad D(\theta) = \theta,$$

an operatorial equation, where $\lambda \in \mathbb{R}$, $D : E \rightarrow E$, $x, y \in E$, and θ is the null element of the space E .

In order to solve equation (1) we consider the following iterative process:

$$(2) \quad x_{n+1} = \lambda D(x_n) + y, \quad n = 0, 1, \dots, x_0 = y.$$

We denote by $S = \{x \in E : \|x\| \leq \rho\}$ the ball of radius ρ and center θ .

Regarding the convergence of iterations (2) we shall consider the following theorem

Theorem 1. *If the application D and the element y from equation (1) verify the following conditions:*

- i. $\|D(x_1) - D(x_2)\| \leq C(\rho) \|x_1 - x_2\|$ for all $x_1, x_2 \in S(\theta)$, where $C : (0, +\infty) \rightarrow (0, +\infty)$ is a functional;
- ii. $\gamma = |\lambda| C(\rho) < 1$;
- iii. $\|y\| \leq (1 - \gamma)\rho$,

then equation (1) admits a unique solution $\bar{x} \in S(\theta, \rho)$.

This solution is obtained as the limit of the sequence $(x_n)_{n=0}^{\infty}$ generated by method (2), and the following estimation holds:

$$(3) \quad \|\bar{x} - x_n\| \leq \frac{\gamma^{n+1}}{1-\gamma} \|y\|, \quad n = 0, 1, \dots$$

Let $D_\varepsilon : E \rightarrow E$ be an application which verifies the conditions:

- i₁. $\|D(x) - D_\varepsilon(x)\| \leq \eta_1(\varepsilon, \rho), \forall x \in B(\theta, \rho)$, where $\eta_1 : [0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$ and $\lim_{\varepsilon \rightarrow 0} \eta_1(\varepsilon, \rho) = 0, \forall \rho > 0$
- ii₁. $\|D_\varepsilon(x_1) - D_\varepsilon(x_2)\| \leq C_\varepsilon(\rho) \|x_1 - x_2\|$ for every $x_1, x_2 \in S(\theta, \rho)$ where $C_\varepsilon : (0, +\infty) \rightarrow (0, +\infty)$;
- iii₁. $|C(\rho) - C_\varepsilon(\rho)| < \eta_2(\varepsilon)$ where $\eta_2 : [0, +\infty) \rightarrow [0, +\infty)$ and $\lim_{\varepsilon \rightarrow 0} \eta_2(\varepsilon) = 0$;
- iv₁. We consider an element for which $\|y - y_\varepsilon\| \leq \eta_3(\varepsilon)$ where $\eta_3 : [0, +\infty) \rightarrow [0, +\infty)$ and $\lim_{\varepsilon \rightarrow 0} \eta_3(\varepsilon) = 0$.

In order to solve equation (1) we consider, instead of iterative method (2) the following iterative procedure:

$$(4) \quad \xi_{n+1} = \lambda D_\varepsilon(\xi_n) + y_\varepsilon, \quad n = 0, 1, \dots, \xi_0 = y_\varepsilon.$$

Regarding the convergence of method (4) we obtain the following theorem:

Theorem 2. *If the conditions of Theorem 1 are fulfilled, the operator D_ε and the element y_ε verify conditions i_1 - iv_1 , and if*

$$(5) \quad \delta = (1 - \gamma) \rho - y \|y\| > 0,$$

then there exists an $\bar{\varepsilon} > 0$, so that for all $\varepsilon < \bar{\varepsilon}$ we have

$$(6) \quad \gamma_\varepsilon = |\lambda| C_\varepsilon(\rho) \leq \gamma + |\lambda| \eta_2(\varepsilon) < 1;$$

and

$$(7) \quad \|y_\varepsilon\| \leq (1 - \gamma_\varepsilon) \rho.$$

Proof. Indeed, from the fact that $\gamma = |\lambda| C(\rho) < 1$ and $\lim_{\varepsilon \rightarrow 0} \eta_2(\varepsilon) = 0$ it follows that there exists a number $\bar{\varepsilon}_1 > 0$ so that for $\varepsilon < \bar{\varepsilon}_1$ we have

$$\gamma_\varepsilon = |\lambda| C_\varepsilon(\rho) \leq |\lambda| C(\rho) + |\lambda| \eta_2(\varepsilon) < 1$$

and

$$\begin{aligned} \|y_\varepsilon\| &\leq \|y\| + \eta_3(\varepsilon) \leq (1 - \gamma) \rho - \delta + \eta_3(\varepsilon) \\ &\leq (1 - \gamma_\varepsilon) \rho + |\lambda| \eta_2(\varepsilon) \rho + \eta_3(\varepsilon) - \delta \leq (1 - \gamma_\varepsilon) \rho \end{aligned}$$

for $\varepsilon < \bar{\varepsilon}_2$ because $\eta_3(\varepsilon) \rightarrow 0$ for $\varepsilon \rightarrow 0$ and $\eta_2(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

If we take now

$$\bar{\varepsilon} = \min\{\bar{\varepsilon}_1, \bar{\varepsilon}_2\}$$

then the theorem is proved. \square

Relations (6) and (7) assure the convergence of sequence $(\xi_n)_{n=0}^\infty$ determined by method (4).

We show now that in the conditions of Theorem 2 we have the following estimation:

$$(8) \quad \|\bar{x} - \xi_n\| \leq \frac{|\lambda| C(\rho) \|\xi_n - \xi_{n-1}\| + |\lambda| \eta_1(\varepsilon, \rho) + \eta_3(\varepsilon)}{1 - \gamma}$$

Indeed, we have

$$\begin{aligned}\|\bar{x} - \xi_n\| &\leq |\lambda| \|D(\bar{x}) - D_\varepsilon(\xi_{n-1})\| + \|y - y_\varepsilon\| \\ &\leq |\lambda| C(\rho) \|\bar{x} - \xi_n\| + |\lambda| C(\rho) \|\xi_n - \xi_{n-1}\| + |\lambda| \eta_1(\varepsilon, \rho) + \eta_3(\varepsilon)\end{aligned}$$

so inequality (8) follows. From (8) it results

$$\|\bar{x} - \bar{\xi}\| \leq \frac{|\lambda| \eta_1(\varepsilon, \rho) + \eta_3(\varepsilon)}{1 - \gamma}, \quad \text{as } n \rightarrow \infty,$$

where $\bar{\xi} = \lim_{n \rightarrow \infty} \xi_n$.

REFERENCES

- [1] Babici, D.M., Ivanov, V.N., *Oțenca polnoi progresnosti prireshenia nelineinyh operatornyh uravnenii metodov prostei iterații*. Jurnal vycislitelnoi matematiki i matematiceskoi fiziki 7, 5 (1967), 988–1000.
- [2] Păvăloiu, I., *Introduction in the Theory of Approximation the Solutions of Equations*, Ed. Dacia 1976 (in Romanian).
- [3] Urabe, M., *Error estimation in numerical solution of equations by iteration process*, J. Sci. Hiroshima Univ. Ser. A-I, 26 (1962), 77–91.

available soon,
← refresh and click here